


## WARNING

KEEP OUT OF REACH OF CHILDREN

For Animal Treatment Only



# MECTIN DRENCH FOR SHEEP

A broad spectrum sheep drench for the control of susceptible gastrointestinal roundworms and lungworms in sheep and lambs including benzimidazole and levamisole resistant strains. Also contains minerals to provide supplementary nutritional support.

## INGREDIENTS:

Abamectin 1 mg/mL, Selenium 0.5 mg/mL, Cobalt 0.4 mg/mL

Copper 2 mg/mL, Zinc 0.6 mg/mL, Iodine 0.8 mg/mL

NET CONTENTS

20 Litre



# MECTIN DRENCH FOR SHEEP



## WARNING:

Dangerous to the Environment. Very toxic to aquatic organisms and terrestrial invertebrates and harmful to the soil environment. Avoid release into the environment or contamination of any water supply with product or empty container.



## PLEASE READ LABEL COMPLETELY BEFORE USE

## HANDLING PRECAUTIONS:

May be harmful if swallowed. Handle with care avoiding skin and eye contact. May cause mild skin irritation. May cause eye irritation.

Repeated exposure may cause skin allergy. Possibly may affect development and/or reproduction. May have effects on/or via lactation.



## PRECAUTIONS:

Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do Not exceed stated dose or dose more frequently than three weekly. In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species. Do not over dose lambs under 15 kg or four weeks of age.

## FIRST AID:

If swallowed, Do NOT induce vomiting. For advice, contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately. If skin or hair contact occurs, remove contaminated clothing and wash skin thoroughly. If splashed in eyes, flush immediately with copious amounts of water.

## INDICATIONS

Contains abamectin 1 mg/mL (a member of the macrocyclic lactone family), selenium 0.5 mg/mL, cobalt 0.4 mg/mL, copper 2 mg/mL, zinc 0.6 mg/mL, iodine 0.8 mg/mL. For the treatment of the following susceptible immature and mature (\*includes inhibited L4) gastrointestinal roundworms and lungworms, including benzimidazole and levamisole resistant strains: *Haemonchus spp.\**, *Teladorsagia (Ostertagia) spp.\**, *Cooperia spp.\**, *Trichostrongylus spp.*, *Oesophagostomum spp.* (adults only) *Nematodirus spp.* (adults only), *Dictyocaulus spp.*

## RESISTANCE MANAGEMENT

Resistance can develop to any anthelmintic. Ask your local animal health advisor or veterinary practitioner for recommended parasite management practices for your area to reduce the development of resistance. It is advised that a resistance test be carried out regularly when using any parasite treatment.

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.**

## DIRECTIONS FOR USE

Shake well before use. Mectin Drench for Sheep is a ready to use oral anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose. Therefore, before treatment, weigh a representative sample of sheep to determine the weight of the heaviest animal. Where there is a large variation in liveweights, drafting into two or more lines may be appropriate before weighing and treating, to avoid under and over dosing. In this situation it is important to choose the dose rate based on the label recommendation for each weight band. Use correct drenching techniques and check accuracy of drenching equipment regularly. Wash out drench gun thoroughly with water before and after use.

## Dose Chart

Do Not overdose lambs under 15 kg or 4 weeks of age.

Dose rate sheep: 1 mL/5kg (equivalent to 0.2mg abamectin/kg, 0.1mg selenium /kg, 0.08mg cobalt /kg, 0.4mg copper /kg, 0.1mg zinc /kg, 0.16mg iodine /kg).

ANIMAL WEIGHT	DOSE	NO. OF DOSES PER 20L PACK
15kg	3 mL	6666
16 - 20 kg	4 mL	5000
21 - 25 kg	5 mL	4000
26 - 30 kg	6 mL	3333
31 - 35 kg	7 mL	2857
36 - 40 kg	8 mL	2500
41 - 45 kg	9 mL	2222
46 - 50 kg	10 mL	2000
51 - 55 kg	11 mL	1818
56 - 60 kg	12 mL	1666
61 - 65 kg	13 mL	1538
66 - 70 kg	14 mL	1428

Over 70 kg dose at the rate of 1 mL/5 kg.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

## WITHHOLDING PERIODS

**Meat:** Sheep producing meat or offal for sale for human consumption must not be sold for slaughter either during treatment or within **21 days** of the last treatment.

**Milk:** Milk intended for sale for human consumption must be discarded during treatment and for not less than **35 days** following the last treatment.

## STORAGE:

Store in original container, tightly closed, in cool, dry conditions. Protect from light. Avoid temperatures below 0° and above 30°C.

## DISPOSAL:

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

## EMERGENCY RESPONSE:

In case of spillage wear appropriate protective clothing. Prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.

Registered pursuant to the ACVM Act 1997, No. A9920.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.  
Registered to: Seneca Holdings Ltd  
Distributed by: Animal Health Direct Ltd  
1229 Maraekakaho Road, Longlands, Hastings,  
New Zealand  
sales@ahdLtd.co.nz. Ph (06) 873 3611

Batch No.:

Expiry date:



9 417726 701271

LMSD20B/04