

HANDLING PRECAUTIONS

Harmful if swallowed, inhaled or absorbed through the skin. May cause eye irritation. Repeated exposure may cause skin allergy. Suspected of damaging fertility or the unborn child and may cause harm to breast-fed children. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. If medical advice is needed, have product container or label at hand. Contains 200g/L benzyl alcohol.

Wear protective gloves and eye protection. Wash hands and exposed skin before meals and after use. Do not inhale vapour. Avoid contact during pregnancy/while nursing. Do not eat, drink or smoke when using this product.

FIRST AID

If swallowed do NOT induce vomiting. If on skin wash thoroughly with plenty of soap and water. If in eyes rinse continuously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin or eye irritation persists or if you feel unwell contact the National Poisons centre 0800 POISON (0800 764 766) or a doctor immediately.

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life and terrestrial invertebrates. Avoid contamination of any water supply with product or empty container. Toxic to the soil environment and harmful to terrestrial vertebrates. Avoid release to the environment. Collect spillage.

EMERGENCY RESPONSE

For specialist advice in an emergency call your local animal health advisor.

BOVIPOR™ ABAMECTIN POUR-ON FOR CATTLE AND DEER

Contains 10mg/mL Abamectin, a member of the macrocyclic lactone family. It is effective against internal and external parasites sensitive to this family. Resistance can develop to any anthelmintic. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

INDICATIONS

Cattle: For the treatment and control of the following mature and immature (*includes inhibited L4) roundworm, lungworms and lice in cattle:

Gastrointestinal worms: Haemonchus spp*, Ostertagia spp*, Cooperia spp*, Trichostrongylus spp; Mature Oesophagostomum spp; Nematodirus spp; Capillaria spp; Trichuris spp; Chabertia spp.

Lungworm: Dictyocaulus spp.

Lice: treatment of biting lice and control of sucking lice.

The efficacy of BoviPor™ is not adversely affected if rain occurs following application. Temporary hair loss occasionally occurs following application therefore do not treat cattle prior to showing. Increases milk production in grazing dairy cows. Data has shown that milk production increases by a median of approximately +0.6kg liquid milk per day when cows or heifers infected with gastrointestinal nematode parasites are treated with an effective anthelmintic; this increase occurs without any reduction in milk quality.

PERSISTENT ACTIVITY

At recommended dose rates BoviPor™ has extended activity against Cooperia and Ostertagia for at least 14 days in cattle.

Deer: For the treatment and control of the following mature roundworm, lungworm and lice in deer:

Gastrointestinal worms: Ostertagia type nematodes, Trichostrongylus axei, Oesophagostomum spp., and Cooperia curticei.

Lungworm: Dictyocaulus spp.

Lice: treatment and control of biting lice.

DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. Shake well before use. BoviPor™ is ready to use through standard pour-on equipment. Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment. Dose rate to be based on the heaviest animal in each group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. Correct drenching technique should be used.

Check the accuracy of drench guns regularly. Do not treat calves under 16 weeks of age.

DOSAGE GUIDE

Cattle and deer: 1mL/20kg bodyweight (500 µg Abamectin/kg bodyweight). Apply along the back line from between the shoulders to the tail using the applicator.

Bodyweight	Dose	5L Pack Treats
120kg	6.0mL	833
150kg	7.5mL	666
200kg	10.0mL	500
250kg	12.5mL	400
300kg	15.0mL	333
350kg	17.5mL	285
400kg	20.0mL	250
450kg	22.5mL	222
500kg	25.0mL	200
550kg	27.5mL	181

Over 550kg use 1mL/20kg bodyweight (2.5mL/50kg bodyweight).

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

MILK: Nil

MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter

either during treatment or:

Cattle within 35 days of cessation of the last treatment

Deer within 28 days of cessation of the last treatment

Bobby calves from cows within 16 days of cessation of the last treatment

STORAGE: Store locked up below 30°C in a cool place, away from direct sunlight. Store in carton, with top secured.

DISPOSAL:

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse empty container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose. This backpack has been designed to be used in an upright position

- 1.Remove the transit cap.
 - 2.Replace with breather cap and draw-off tube.
 - 3.Attach applicator gun to breather cap.
- ENSURE THAT YOU USE THE RECOMMENDED APPLICATOR GUN.
4. **Always use or carry the pack in an upright position.** Optional backpack straps are provided.
 - 5.At conclusion of use replace transit cap.

Registered pursuant to the ACVM Act 1997, No. A11132
See www.foodsafety.govt.nz for registration conditions

Registered to WSD Agribusiness Pty Ltd
NZ Agent Orion AgriScience Ltd
Unit 1, 15 Sir Gil Simpson Drive
Harewood, Christchurch 8053

Batch

Expires

orion
AGRISCIENCE

